

Novocastra™ Ready-to-Use Mouse Monoclonal Antibody Endothelial Cell Marker (CD34)

Product Code: RTU-END

Intended Use	FOR RESEARCH USE ONLY.
Specificity	Human endothelial cells (CD34 Antigen)
Clone	QBEnd/10
Ig Class	IgG1
Antigen Used for Immunizations	Detergent solubilized vesicular suspension prepared from a perfusate of human term placenta.
Hybridoma Partner	Mouse myeloma (p3/NS-0).
Preparation	Tissue culture supernatant diluted in 5% horse serum in PBS containing 12 mM sodium azide. Volume as indicated on vial label.
Effective on Frozen Tissue	Yes
Effective on Paraffin Wax Embedded Tissue	Yes, trypsin digestion of paraffin sections is recommended in preference to high temperature antigen unmasking techniques. It produces stronger staining and is effective in reducing background elastic staining.
Recommendations on Use	<p>Immunohistochemistry: Typical working dilution: NEAT. 15 minutes primary antibody incubation at 25 °C when used in conjunction with the Novostain Universal Detection Kit (Ready to Use), code NCL-RTU-D. Recommendations on use will differ if other detection systems are used eg Standard ABC technique. Western Blotting: Not recommended.</p> <p>Not recommended for use on Ventana automated staining systems (Ventana Medical Systems Inc., USA).</p>
Positive Controls	Immunohistochemistry: Any tissue in which endothelial cells are readily identified, for example, normal human skin.
Staining Pattern	Endothelial cell cytoplasm.
Storage and Stability	Store ready-to-use prediluted liquid antibody at 4 °C. Return to 4 °C immediately after use. Under these conditions, there is no significant loss in product performance up to the expiry date indicated on the vial label.
General Overview	CD34 antigen is a single chain transmembrane glycoprotein with a molecular weight of 110 kD. The CD34 antigen is selectively expressed on human lymphoid and myeloid hemopoietic progenitor cells. The antigen is also expressed on vascular endothelium.
General References	<p>Watanabe T, Oda Y, Tamiya S, et al.. Journal of Clinical Pathology. 54: 631–636 (2001).</p> <p>Cox G, Walker R A, Andi A, et al.. Lung Cancer. 29 (3): 169–177 (2000).</p> <p>Anthony P P and Ramani P. Journal of Clinical Pathology. 44: 29–32 (1991).</p> <p>Dhillon A P, Sankey E A and More L. Journal of Pathology. 162: 274 (1990).</p> <p>Fletcher C D M and Ramani P. Journal of Pathology. 162: 273 (1990).</p> <p>Ramani P, Bradley N J and Fletcher C D M. Histopathology. 17: 237–242 (1990).</p> <p>Sankey E A, More L and Dhillon A P. Journal of Pathology. 161: 267–271 (1990).</p>



Instructions for Use

Trypsin Digestion for Immunohistochemical Demonstration on Paraffin Sections

1. Preheat the following to 37 °C using a water bath:
 - (i) 200 mL of TBS
 - (ii) 200 mL of distilled water.
2. Dissolve 0.2 g Trypsin 250 and 0.2 g Calcium chloride in the 200 mL of TBS.
3. Once the Trypsin solution is at 37 °C, pH to 7.8 with 1 M sodium hydroxide.
4. Place rehydrated paraffin sections in the distilled water to preheat the sections to 37 °C for a minimum of 5 minutes.
5. Incubate sections in Trypsin solution at 37 °C. The time required will depend on the antibody and tissue, however, 30 minutes is usually sufficient.
6. Rinse sections in running tap water.
7. Proceed with immunohistochemistry protocol.

Reagents Required but not Supplied

50 mM Tris-buffered saline

Trypsin 250: Difco order code 0152-13 (available from Becton Dickinson).

Calcium chloride

1 M Sodium Hydroxide

** Trypsin containing chymotrypsin should always be used. The enzyme activities can vary from a supplier and between batches. Such variations may affect the incubation time required.*